arXiv:1404.6600v1 [math.CV] 26 Apr 2014

ON AN INEQUALITY CONCERNING THE POLAR DERIVATIVE OF A POLYNOMIAL WITH RESTRICTED ZEROS

N. A. RATHER AND SUHAIL GULZAR

Abstract. Let $D_\alpha P(z) = nP(z) + (\alpha - z)P'(z)$ denote the polar derivative of a polynomial $P(z)$ of degree n with respect to a point $\alpha \in \mathbb{C}$. In this paper, we present a correct proof, independent of Laguerre's theorem, of an inequality concerning the polar derivative of a polynomial with restricted zeros recently formulated by K. K. Dewan, Naresh Singh, Abdullah Mir, [Extensions of some polynomial inequalities to the polar derivative, *J. Math. Anal. Appl.,* **352** (2009) 807-815].

1. Introduction

If $P(z)$ is a polynomial of degree n having all its zeros in $|z| \leq 1$, it was shown by Turán [9] that

$$\max_{|z|=1} \left|P'(z)\right| \geq \frac{n}{2} \max_{|z|=1} |P(z)|. \tag{1.1}$$

The inequality (1.1) becomes equality for $P(z) = (z+1)^n$.

As an extension of (1.1), Malik [6] proved that if all the zeros of polynomial $P(z)$ of degree n lie in $|z| \leq k$, $k \leq 1$, then

$$\max_{|z|=1} \left|P'(z)\right| \geq \frac{n}{1+k} \max_{|z|=1} |P(z)|. \tag{1.2}$$

By considering the class of polynomials $P(z) = a_n z^n + \sum_{j=\mu}^{n} a_{n-j} z^{n-j}$, $1 \leq \mu \leq n$, of degree n having all their zeros in $|z| \leq k$, $k \leq 1$, Aziz and Shah [1] proved

$$\max_{|z|=1} \left|P'(z)\right| \geq \frac{n}{1+k^\mu} \left\{ \max_{|z|=1} |P(z)| + \frac{1}{k^{n-\mu}} \min_{|z|=k} |P(z)| \right\}. \tag{1.3}$$

Let $D_\alpha P(z)$ denote the polar derivative of a polynomial $P(z)$ of degree n with respect to a point $\alpha \in \mathbb{C}$, then (see [7])

$$D_\alpha P(z) = nP(z) + (\alpha - z)P'(z).$$

The polynomial $D_\alpha P(z)$ is of degree at most $n-1$ and it generalizes the ordinary derivative in the sense that

$$\lim_{\alpha \to \infty} \frac{D_\alpha P(z)}{\alpha} = P'(z)$$

uniformly with respect to z for $|z| \leq R$, $R > 0$.

Dewan et al. [4] (see also [8]) extended inequality (1.3) to polar derivative

2010 Mathematics Subject Classification. 30A10, 30C10.

Key words and phrases. Polynomials; Inequalities in the complex domain; Polar derivative; Bernstein's inequality.

and proved that if $P(z) = a_n z^n + \sum_{j=\mu}^{n} a_{n-j} z^{n-j}$, $1 \leq \mu \leq n$, is a polynomial of degree n having all its zeros in $|z| \leq k$, $k \leq 1$, then for every real and complex number α with $|\alpha| \geq k^\mu$,

$$\max_{|z|=1} |D_\alpha P(z)| \geq \frac{n\left(|\alpha| - k^\mu\right)}{1+k^\mu} \max_{|z|=1} |P(z)| + \frac{n\left(|\alpha|+1\right)}{k^{n-\mu}\left(1+k^\mu\right)} \min_{|z|=k} |P(z)|. \quad (1.4)$$

While seeking the desired refinement of inequality (1.4), recently Dewan et al. [3] have made an incomplete attempt by claiming to have proved the following result.

Theorem 1.1. *If $P(z) = a_n z^n + \sum_{j=\mu}^{n} a_{n-j} z^{n-j}$, $1 \leq \mu \leq n$, is a polynomial of degree n having all its zeros in $|z| \leq k$, $k \leq 1$, then for every real or complex number α with $|\alpha| \geq k^\mu$,*

$$\begin{aligned} \max_{|z|=1} |D_\alpha P(z)| \geq & \frac{n\left(|\alpha| - k^\mu\right)}{1+k^\mu} \max_{|z|=1} |P(z)| + \frac{n\left(|\alpha|+1\right)}{k^{n-\mu}\left(1+k^\mu\right)} m \\ & + n\left(\frac{k^\mu - A_\mu}{1+k^\mu}\right) \max_{|z|=1} |P(z)| + \frac{n(A_\mu - k^\mu)}{k^n(1+k^\mu)} m \end{aligned} \quad (1.5)$$

where $m = \min_{|z|=k} |P(z)|$ and

$$A_\mu = \frac{n\left(|a_n| - \frac{m}{k^n}\right) k^{2\mu} + \mu |a_{n-\mu}| k^{\mu-1}}{n\left(|a_n| - \frac{m}{k^n}\right) k^{\mu-1} + \mu |a_{n-\mu}|}. \quad (1.6)$$

The proof of Theorem 1.1 given by Dewan et al. [3] is not correct. The reason being that the authors in [3], deduce by using Lemma 7[3] between the lines 8 to 10, on page 814, that if $F(z) = P(z) - m\lambda z^n/k^n$ has all its zeros in $|z| < k, k \leq 1$, then for every $\alpha \in \mathbb{C}$ with $|\alpha| \geq k^\mu$, $1 \leq \mu \leq n$ all the zeros of the polar derivative $D_\alpha F(z)$ also lie in $|z| < k$ which need not be true in general for $1 \leq \mu \leq n$. Here, Lemma 7 [3] is applicable only when $\mu = 1$. For $1 < \mu \leq n$ the assertion is not true, since for $k < 1$, $k^\mu < k$.

The immediate counterexample $P(z) = 4z^2 - 1$, $\mu = 2$ having all its zeros in $|z| < k = 3/5 < 1$ demonstrates, by taking $\alpha = 2/5 > k^\mu$ that the zero of $D_\alpha P(z) = \frac{16z}{5} - 2$ lie in $|z| > k = 3/5$.

The main aim of this paper is to present a correct proof, independent of Laguerre's theorem, of Theorem 1.1.

2. Lemmas

For the proof of above theorem, we need the following lemmas.

Lemma 2.1. *If $P(z) = a_n z^n + \sum_{j=\mu}^{n} a_{n-j} z^{n-j}$, $1 \leq \mu \leq n$, is a polynomial of degree n having all its zeros in $|z| \leq k$, $k \leq 1$ and $Q(z) = z^n \overline{P(1/\overline{z})}$, then for $|z| = 1$*

$$|Q'(z)| \leq S_\mu |P'(z)| \ \left(\leq k^\mu |P'(z)|\right) \quad (2.1)$$

where

$$S_\mu = \frac{n|a_n| k^{2\mu} + \mu |a_{n-\mu}| k^{\mu-1}}{n|a_n| k^{\mu-1} + \mu |a_{n-\mu}|} \quad (2.2)$$

and

$$\frac{\mu}{n} \left|\frac{a_{n-\mu}}{a_n}\right| \leq k^\mu.$$

The above lemma is due to Aziz and Rather [2].

Lemma 2.2. *Let $P(z) = a_n z^n + \sum_{j=\mu}^{n} a_{n-j} z^{n-j}$, $1 \leq \mu \leq n$, be a polynomial of degree n having all its zeros in $|z| \leq k$, $k \leq 1$, then for every real or complex number α with $|\alpha| \geq S_\mu$,*

$$|D_\alpha P(z)| \geq n \left(\frac{|\alpha| - S_\mu}{1 + k^\mu} \right) |P(z)| \qquad for \quad |z| = 1. \tag{2.3}$$

Proof. If $Q(z) = z^n \overline{P(1/\overline{z})}$, then $P(z) = z^n \overline{Q(1/\overline{z})}$ and it can be easily verified that for $|z| = 1$

$$\begin{aligned} |Q'(z)| &= |nP(z) - zP'(z)| \\ &\geq |nP(z)| - |zP'(z)|. \end{aligned}$$

Equivalently, we have

$$|Q'(z)| + |P'(z)| \geq n|P(z)| \quad \text{for} \quad |z| = 1. \tag{2.4}$$

This implies with the help of Lemma 2.1 and inequality (2.4), that

$$\begin{aligned} (1 + k^\mu) |P'(z)| &= |P'(z)| + k^\mu |P'(z)| \\ &\geq |P'(z)| + |Q'(z)| \\ &\geq n|P(z)|, \end{aligned}$$

which implies

$$\left|P'(z)\right| \geq \frac{n}{1 + k^\mu} |P(z)| \quad \text{for} \quad |z| = 1. \tag{2.5}$$

Now, for every real or complex number α with $|\alpha| \geq S_\mu$,

$$\begin{aligned} |D_\alpha P(z)| &= |nP(z) + (\alpha - z)P'(z)| \\ &\geq |\alpha||P'(z)| - |nP(z) - zP'(z)| \\ &= |\alpha||P'(z)| - |Q'(z)| \qquad \text{for} \quad |z| = 1. \end{aligned} \tag{2.6}$$

Combing inequalities (2.5), (2.6) and Lemma 2.1, we get for $|z| = 1$,

$$\begin{aligned} |D_\alpha P(z)| &\geq (|\alpha| - S_\mu) |P'(z)| \\ &\geq n \left(\frac{|\alpha| - S_\mu}{1 + k^\mu} \right) |P(z)| \end{aligned}$$

This proves Lemma 2.2. □

Lemma 2.3. *If $P(z) = \sum_{j=1}^{n} a_j z^j$ is a polynomial of degree n having all its zeros in $|z| \leq k$, $k \leq 1$ and $m = \min_{|z|=k} |P(z)|$, then*

$$\max_{|z|=1} |P(z)| \geq \frac{m}{k^n} \tag{2.7}$$

and in particular,

$$|a_n| > \frac{m}{k^n}. \tag{2.8}$$

Proof. Since the polynomial $P(z)$ has all its zeros in $|z| \leq k$, $k \leq 1$, the polynomial $Q(z) = z^n \overline{P(1/\overline{z})}$ has no zero in $|z| < 1/k$, $1/k \geq 1$. We assume that $Q(z)$ has no zero on $|z| = 1/k$, for otherwise the result holds trivially. Since $Q(z)$ has no zeros in $|z| \leq 1/k$, by Minimum Modulus Principle

$$|Q(z)| \geq \min_{|z|=1/k} |Q(z)| \quad \text{for} \quad |z| \leq 1/k \ \text{ where } \ 1/k \geq 1,$$

which in particular gives,

$$|Q(z)| \geq \frac{1}{k^n}\min_{|z|=k}|P(z)| \quad \text{for} \quad |z| \leq 1 \quad \text{and} \quad |a_n| = |Q(0)| > \frac{1}{k^n}\min_{|z|=k}|P(z)|.$$

That is,

$$\max_{|z|=1}|P(z)| = \max_{|z|=1}|Q(z)| \geq \frac{m}{k^n} \quad \text{and} \quad |a_n| > \frac{m}{k^n}.$$

This completes the proof of Lemma 2.3.. □

Lemma 2.4. *The function*

$$S_\mu(x) = \frac{nxk^{2\mu} + \mu|a_{n-\mu}|k^{\mu-1}}{nxk^{\mu-1} + \mu|a_{n-\mu}|}. \tag{2.9}$$

where $k \leq 1$ and $\mu \geq 1$, is a non-increasing function of x.

Proof. The proof follows by considering the first derivative test for $S_\mu(x)$. □

Lemma 2.5. *If $P(z) = a_n z^n + \sum_{\nu=\mu}^{n} a_{n-\nu} z^{n-\nu}$, $1 \leq \mu \leq n$, is a polynomial of degree n, having all its zeros in $|z| \leq k$, $k \leq 1$, then for every $\alpha \in \mathbb{C}$ with $|\alpha| \geq k^\mu$,*

$$|D_\alpha P(z)| \geq \frac{|\alpha|mn}{k^n} \qquad \text{for} \qquad |z| = 1 \tag{2.10}$$

where $m = \min_{|z|=k}|P(z)|$.

Proof. By hypothesis all the zeros of polynomial $P(z) = a_n z^n + \sum_{\nu=\mu}^{n} a_{\nu-\mu} z^{\nu-\mu}$, $1 \leq \mu \leq n$, lie in $|z| \leq k$, $k \leq 1$. If $P(z)$ has a zero on $|z| = k$ then $m = 0$ and the result holds trivially. Hence, we suppose that all the zeros of $P(z)$ lie in $|z| < k$, $k \leq 1$, so that $m > 0$ and $m \leq |P(z)|$ for $|z| = k$, mow if λ is any complex number such that $|\lambda| < 1$, then

$$\left|\frac{m\lambda z^n}{k^n}\right| < |P(z)| \qquad \text{for} \qquad |z| = k.$$

Since all the zeros of $P(z)$ lie in $|z| < k$, it follows by Rouche's theorem that all the zeros of

$$F(z) = P(z) - \frac{m\lambda z^n}{k^n}$$

also lie in $|z| < k, k \leq 1$. Applying Lemma 2.1 to the polynomial $F(z)$, we get

$$k^\mu |F'(z)| \geq |G'(z)| \qquad \text{for} \qquad |z| = 1.$$

where $G(z) = z^n\overline{F(1/\overline{z})} = z^n\overline{P(1/\overline{z})} + \frac{m\overline{\lambda}}{k^n} = Q(z) + + \frac{m\overline{\lambda}}{k^n}$. Equivalently for $|z| = 1$, we have

$$k^\mu \left|P'(z) - \frac{mn\lambda z^{n-1}}{k^n}\right| \geq |Q'(z)|. \tag{2.11}$$

Moreover, by Gauss-Lucas theorem that all the zeros of polynomial $F' = P'(z) - mn\lambda z^{n-1}/k^n$ lie in $|z| < k$, $k \leq 1$ for every λ with $|\lambda| < 1$. This implies

$$|P'(z)| \geq mn|z|^{n-1}/k^n \quad \text{for} \quad |z| \geq k. \tag{2.12}$$

Now, choosing the argument of λ in the left hand side of (2.11) such that

$$k^{\mu}\left|P'(z)-\frac{mn\lambda z^{n-1}}{k^n}\right| = k^{\mu}\left\{|P'(z)|-\frac{|\lambda|mn}{k^n}\right\} \quad \text{for} \quad |z|=1,$$

which is possible by (2.12), we get

$$k^{\mu}|P'(z)|-\frac{|\lambda|mn}{k^{n-\mu}} \geq |Q'(z)| \quad \text{for} \quad |z|=1.$$

Letting $|\lambda| \to 1$, we obtain

$$k^{\mu}|P'(z)|-\frac{mn}{k^{n-\mu}} \geq |Q'(z)| \quad \text{for} \quad |z|=1. \tag{2.13}$$

Again, for $|z|=1$ and $\alpha \in \mathbb{C}$, we have

$$\begin{aligned} |D_{\alpha}P(z)| &= |nP(z)+(\alpha-z)P'(z)| \\ &\geq |\alpha||P'(z)|-|nP(z)-zP'(z)| \\ &= |\alpha||P'(z)|-|Q'(z)| \end{aligned}$$

Combining this with inequality (2.13), we obtain for $|z|=1$ and $\alpha \in \mathbb{C}$ with $|\alpha| \geq k^{\mu}$,

$$|D_{\alpha}P(z)| \geq (|\alpha|-k^{\mu})|P'(z)|+k^{\mu}\frac{mn}{k^n} \tag{2.14}$$

Inequality (2.14) in conjunction with (2.12) gives for $|z|=1$ and $|\alpha| \geq k^{\mu}$,

$$\begin{aligned} |D_{\alpha}P(z)| &\geq (|\alpha|-k^{\mu})\frac{mn}{k^n}+k^{\mu}\frac{mn}{k^n} \\ &= \frac{|\alpha|mn}{k^n}. \end{aligned} \tag{2.15}$$

This completes the proof of Lemma 2.5.

□

The next lemma is due to Gardner, Govil and Weems [5].

Lemma 2.6. *If $P(z)=a_0+\sum_{j=\mu}^{n}a_jz^j$, $1 \leq \mu \leq n$, is a polynomial of degree n having no zero in $|z| \leq k$, $k \geq 1$, then*

$$s_0 \geq k^{\mu} \tag{2.16}$$

where

$$s_0 = k^{\mu+1}\left\{\frac{\left(\frac{\mu}{n}\right)\frac{|a_{\mu}|}{|a_0|-m}k^{\mu-1}+1}{\left(\frac{\mu}{n}\right)\frac{|a_{\mu}|}{|a_0|-m}k^{\mu+1}+1}\right\}.$$

The following Lemma can be easily verified by applying Lemma 2.6 to $Q(z)=z^n\overline{P(1/\overline{z})}$ where $P(z)=a_nz^n+\sum_{j=\mu}^{n}a_{n-j}z^{n-j}$, $1 \leq \mu \leq n$.

Lemma 2.7. *If $P(z)=a_nz^n+\sum_{j=\mu}^{n}a_{n-j}z^{n-j}$, $1 \leq \mu \leq n$, is a polynomial of degree n having all its zeros in $|z| \leq k$, $k \leq 1$, then*

$$A_{\mu} \leq k^{\mu} \tag{2.17}$$

where A_{μ} is defined in (1.6).

3. Proof of theorem

Proof of theorem 1.1. Proceeding similarly as in the proof of lemma 2.5, the polynomial $P(z) - \frac{m\lambda z^n}{k^n}$ has all its zeros in $|z| \leq k$, $k \leq 1$, for every real or complex λ with $|\lambda| < 1$. Applying Lemma 2.2 to the polynomial $P(z) - \frac{m\lambda z^n}{k^n}$, we obtain for $|\alpha| \geq S'_\mu$,

$$\left|D_\alpha\Big\{P(z) - \frac{m\lambda z^n}{k^n}\Big\}\right| \geq n\left(\frac{|\alpha| - S'_\mu}{1+k^\mu}\right)\left|P(z) - \frac{m\lambda z^n}{k^n}\right| \quad \text{for} \quad |z| = 1$$

where

$$S'_\mu = \frac{n\left|a_n - \frac{m\lambda}{k^n}\right| k^{2\mu} + \mu|a_{n-\mu}|k^{\mu-1}}{n\left|a_n - \frac{m\lambda}{k^n}\right| k^{\mu-1} + \mu|a_{n-\mu}|}. \tag{3.1}$$

That is,

$$\left|D_\alpha P(z) - \frac{\lambda mn\alpha z^{n-1}}{k^n}\right| \geq n\left(\frac{|\alpha| - S'_\mu}{1+k^\mu}\right)\left|P(z) - \frac{m\lambda z^n}{k^n}\right| \quad \text{for} \quad |z| = 1 \tag{3.2}$$

Since for every λ with $|\lambda| < 1$, we have by inequality (2.8) of Lemma 2.7,

$$\left|a_n - \frac{m\lambda}{k^n}\right| \geq |a_n| - \frac{m|\lambda|}{k^n} \geq |a_n| - \frac{m}{k^n} \tag{3.3}$$

therefore, it follows by Lemma 2.4 that for every λ with $|\lambda| < 1$,

$$\begin{aligned} S'_\mu =& \frac{n\left|a_n - \frac{m\lambda}{k^n}\right| k^{2\mu} + \mu|a_{n-\mu}|k^{\mu-1}}{n\left|a_n - \frac{m\lambda}{k^n}\right| k^{\mu-1} + \mu|a_{n-\mu}|} \\ \leq& \frac{n\left(|a_n| - \frac{m}{k^n}\right) k^{2\mu} + \mu|a_{n-\mu}|k^{\mu-1}}{n\left(|a_n| - \frac{m}{k^n}\right) k^{\mu-1} + \mu|a_{n-\mu}|} = A_\mu \end{aligned} \tag{3.4}$$

Using (3.4) in (3.2), we obtain for $|\alpha| \geq A_\mu (\geq S'_\mu)$,

$$\left|D_\alpha P(z) - \frac{\lambda mn\alpha z^{n-1}}{k^n}\right| \geq n\left(\frac{|\alpha| - A_\mu}{1+k^\mu}\right)\left|P(z) - \frac{m\lambda z^n}{k^n}\right| \quad \text{for} \quad |z| = 1. \tag{3.5}$$

Now, choosing the argument of λ in the right hand side of inequality (3.5) such that

$$\left|D_\alpha P(z) - \frac{\lambda mn\alpha z^{n-1}}{k^n}\right| = |D_\alpha P(z)| - \frac{|\lambda||\alpha|mnz^{n-1}}{k^n} \tag{3.6}$$

which is possible by Lemma 2.5, we obtain by using Lemma 2.7 for $|\alpha| \geq k^\mu (\geq A_\mu)$ and $|z| = 1$,

$$|D_\alpha P(z)| - \frac{|\lambda||\alpha|mn|z|^{n-1}}{k^n} \geq n\left(\frac{|\alpha| - A_\mu}{1+k^\mu}\right)\left\{|P(z)| - \frac{|\lambda|m|z|^n}{k^n}\right\},$$

equivalently,

$$|D_\alpha P(z)| \geq n\left(\frac{|\alpha| - A_\mu}{1+k^\mu}\right)|P(z)| + |\lambda|\frac{mn}{k^n}\left(\frac{|\alpha|k^\mu + A_\mu}{1+k^\mu}\right). \tag{3.7}$$

Letting $|\lambda| \to 1$, in (3.7), we get

$$|D_\alpha P(z)| \geq n\left(\frac{|\alpha| - A_\mu}{1+k^\mu}\right)|P(z)| + \frac{mn}{k^n}\left(\frac{|\alpha|k^\mu + A_\mu}{1+k^\mu}\right) \quad \text{for} \quad |z| = 1,$$

which implies for every $\alpha \in \mathbb{C}$ with $|\alpha| \geq k^{\mu}$, $1 \leq \mu \leq n$,

$$\max_{|z|=1} |D_{\alpha}P(z)| \geq \frac{n\left(|\alpha| - k^{\mu}\right)}{1+k^{\mu}} \max_{|z|=1} |P(z)| - \frac{n\left(|\alpha|+1\right)}{k^{n-\mu}\left(1+k^{\mu}\right)} m$$
$$+ n\left(\frac{k^{\mu} - A_{\mu}}{1+k^{\mu}}\right) \max_{|z|=1} |P(z)| + \frac{n(A_{\mu} - k^{\mu})}{k^{n}(1+k^{\mu})} m.$$

This completes the proof of Theorem 1.1. □

References

[1] A. Aziz and W. M. Shah, An integral mean estimate for polynomial, *Indian J. Pure Appl. Math.,* **28** (1997) 1413-1419.

[2] A. Aziz and N. A. Rather, Some Zygmund type L^q inequalities for polynomials, *J. Math. Anal. Appl.,* **289** (2004) 14-29.

[3] K. K. Dewan, Naresh Singh, Abdullah Mir, Extensions of some polynomial inequalities to the polar derivative, *J. Math. Anal. Appl.,* **352** (2009) 807-815.

[4] K. K. Dewan, Naresh Singh, Roshan Lal, Inequalities for the polar derivatives of a polynomials, *Int. J. Pure Appl. Math.,* **33** (2006) 109-117.

[5] R.B. Gardner, N.K. Govil, A. Weems, Some results concerning rate of growth of polynomials, *East J. Approx.,* **10** (2004) 301312.

[6] M. A. Malik, On the derivative of a polynomial, *J. London Math. Soc.,* **2** (1969) 5760.

[7] M. Marden, *Geometry of Polynomials*, Math. Surveys No. 3, Amer. Math. Soc. Providence R. I. 1966.

[8] N. A. Rather and M. I. Mir, Some refinements of inequalities for the polar derivative of polynomials with restricted zeros, *Int. J. Pure and Appl. Math.*, **41** (2007), 1065-1074.

[9] P. Turán, Über die Ableitung von Polynomen, *Compositio Mathematica* **7** (1939), 89-95 (German).

Department of Mathematics, University of Kashmir, Srinagar, Hazratbal 190006, India

E-mail address: dr.narather@gmail.com

E-mail address: sgmattoo@gmail.com